

CMK2015-503-119

CMK2015-503-119

LADYANNA x CMK2008-071-009



MEIJER
POTATO

Building on potential

Purpose: French fries

Characteristics:

Maturity	6,0	
Colour of skin	Y	
Colour of flesh	4,7	
Dry matter percentage	20,9	
Suitability for cooking	7,8 AB	
Shape of tuber	LO	
Tuber length (50-60 mm)	8,9 cm	
Average size (mm)	57	
Dormancy	7,2	

Added Value:

White flesh and long shape variety for premium market.

Robust plant, tolerant to climatic stress.

Excellent frying colour throughout the storage period.

Gives high yields from all soil types.

Uniform tuber size in high marketable yields.

Resistances :

Common scab	7	
Bruise index (0-50)	17	
Secondary growth	7	
Yntn-virus	8	
Foliage blight	2	
Fusarium	5	
Ro1	9	
Ro2/3	3	
Ro5		
Pa2	9	
Pa3	9	
wz 1 (D1)	10	
wz 2 (G1)		
wz 6 (O1)	5	
wz 18 (T1)	1	

Planting distance:

Size	Rowspace (cm between plants)		Plants/hectare
	75 cm	90 cm	
28/35	33	28	40404
35/50	41	34	32520
50/55	44	37	30303



CMK2015-503-119




MEIJER 
POTATO
Building on potential

Purpose: French fries




Planting

Wait for eyes waking up before planting.
Do not plant cold seed in cold soil.
Don't de-sprout the seeds before planting.
Plant at a normal depth.



Fertiliser input (kg/ha):

Depending on the soil supply and applicable national legislation; total amount required by the variety.



Nitrogen: (N) 200
Phosphate: (P₂O₅) 110
Potassium: (K₂O) 350

Total
200
110
350

To prevent mineral deficiencies, apply these elements in time (B, Mg, Mn, S)



Crop protection

Susceptible to the use of 450 g/ha of Metribuzin at pre-emergence.
Susceptible to the use of 150 g/ha of Metribuzin at post-emergence.
Variety susceptible to foliar late blight, do not neglect blight control.
Start late blight control in time.



Harvest and storage

Wait for complete seed disintegration before harvesting.
After harvest, start immediately with drying the potatoes.



Storage:
7 °C
9 months

Key advice to succeed this variety:

Wait for eyes waking up before planting.
Susceptible to the use of 450 g/ha of Metribuzin at pre-emergence.
Susceptible to the use of 150 g/ha of Metribuzin at post-emergence.
Variety susceptible to foliar late blight, do not neglect blight control.
After harvest, start immediately with drying the potatoes.