

CMK2012-016-016

CMK2012-016-016

CMK2005-210-007 x LADYANNA



MEIJER
POTATO

Building on potential

Purpose: French fries

Characteristics:

Maturity	5,9	
Colour of skin	Y	
Colour of flesh	6,3	
Dry matter percentage	22,4	
Suitability for cooking	7,9 B	
Shape of tuber	LO	
Tuber length (50-60 mm)	8,6 cm	
Average size (mm)	57	
Dormancy	7,7	

Added Value:

Uniform tuber size in high marketable yields.

Robust plant, tolerant to climatic stress.

Excellent frying colour throughout the storage period.

Gives high yields from all soil types.

Long dormancy suitable for long storage.

Resistances :

Common scab	7	
Bruise index (0-50)	11	
Secondary growth	7	
Yntn-virus	10	
Foliage blight	4	
Fusarium	7	
Ro1	9	
Ro2/3	8	
Ro5		
Pa2	7	
Pa3	1	
wz 1 (D1)	5	
wz 2 (G1)		
wz 6 (O1)	3	
wz 18 (T1)	5	

Planting distance:

Size	Rowspace (cm between plants)		Plants/hectare
	75 cm	90 cm	
28/35	33	28	40404
35/50	41	34	32520
50/55	44	37	30303



CMK2012-016-016




MEIJER 
POTATO
Building on potential

Purpose: French fries




Planting

Wait for eyes waking up before planting.
Do not plant cold seed in cold soil.
Don't de-sprout the seeds before planting.
Plant at a normal depth.
Seems to be have a susceptibility to powdery scab; choose the right soils.



Fertiliser input (kg/ha):

Depending on the soil supply and applicable national legislation; total amount required by the variety.



Nitrogen: (N) 250
Phosphate: (P₂O₅) 110
Potassium: (K₂O) 350

Total
250
110
350

To prevent mineral deficiencies, apply these elements in time (B, Mg, Mn, S)

Variety with low needs of nitrogen.
Too much nitrogen affect the yield in a negativ way.



Crop protection

Susceptible to the use of 150 g/ha of Metribuzin at post-emergence.
Variety susceptible to foliar late blight, do not neglect blight control.
Start late blight control in time.



Harvest and storage

Wait for complete seed disintegration before harvesting.
After harvest, start immediately with drying the potatoes.



Storage:
7 °C
8 months

Key advice to succeed this variety:

Wait for eyes waking up before planting.
Seems to be have a susceptibility to powdery scab; choose the right soils.
Susceptible to the use of 150 g/ha of Metribuzin at post-emergence.
Variety susceptible to foliar late blight, do not neglect blight control.
After harvest, start immediately with drying the potatoes.