

SOUND

CMK2008-622-009

MONDIAL x CMK2003-707-003



MEIJER
POTATO

Building on potential



Purpose: Table

Foliar late blight resistant variety



Characteristics:

Maturity	6,6	
Colour of skin	LY	
Colour of flesh	6,5	
Dry matter percentage	19,8	
Suitability for cooking	7,4 AB	
Shape of tuber	LO	
Tuber length (50-60 mm)	8,8 cm	
Average size (mm)	57	
Dormancy	6,9	

Added Value:

Very suitable for home french fries.

Foliar late blight resistant.

Very nice skin quality.

Gives high yields from all soil types.

Drought tolerant variety.

Resistances :

Common scab	7		
Bruise index (0-50)	18		
Secondary growth	7		
Yntn-virus	7		
Foliage blight	9		
Fusarium	5		
Ro1	9		
Ro2/3	4		wz 1 (D1) 10
Ro5	8		wz 2 (G1)
Pa2	3		wz 6 (O1) 4
Pa3	2		wz 18 (T1) 4

Planting distance:

Size	Rowspace (cm between plants)		Plants/hectare
	75 cm	90 cm	
28/35	30	25	44444
35/55	38	32	35088
55/60	42	35	31746
55/60 CUT	35	29	38095
30/40	32	27	41667
40/50	35	29	38095
50/60	39	33	34188



SOUND




Purpose: Table




Planting

Before planting keep the seed quiet and plant them out of cold storage.
Do not store the seed with Talent (Carvon), which stops the sprouting capacity.
Do not cultivate on soils prone to common scab.
Plant at a normal depth.



Fertiliser input (kg/ha):

Depending on the soil supply and applicable national legislation; total amount required by the variety.



Nitrogen: (N) 200
Phosphate: (P₂O₅) 110
Potassium: (K₂O) 350

Total
200
110
350

To prevent mineral deficiencies, apply these elements in time (B, Mg, Mn, S)

Do not use potassium chloride in order to avoid a low dry matter level.



Crop protection

Susceptible to the use of 450 g/ha of Metribuzin at pre-emergence.
Take care of the late blight foliar resistance, do not neglect your blight program.



Harvest and storage

Wait for a proper set skin before harvest.
Take care to avoid mechanical damages on the skin during the harvest.
Short dormancy; after harvest and wound healing cool down quickly and keep temperature stable.
Start sprouting inhibitor application as soon as potatoes are dry.



Storage:
7 °C
8 months

Key advice to succeed this variety:

Before planting keep the seed quiet and plant them out of cold storage.
Do not use potassium chloride in order to avoid a low dry matter level.
Take care of the late blight foliar resistance, do not neglect your blight program.
Short dormancy; after harvest and wound healing cool down quickly and keep temperature stable.

Don't use
Talent
to store
the seeds